PHYSIOLOGICAL ECOLOGY - ORIGINAL RESEARCH



Tree growth and water-use in hyper-arid *Acacia* occurs during the hottest and driest season

Gidon Winters¹ · Dennis Otieno² · Shabtai Cohen³ · Christina Bogner⁴ · Gideon Ragowloski¹ · Indira Paudel⁵ · Tamir Klein⁵

Received: 10 April 2018 / Accepted: 13 August 2018 © Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Drought-induced tree mortality has been recently increasing and is expected to increase further under warming climate. Conversely, tree species that survive under arid conditions might provide vital information on successful drought resistance strategies. Although *Acacia (Vachellia)* species dominate many of the globe's deserts, little is known about their growth dynamics and water-use in situ. Stem diameter dynamics, leaf phenology, and sap flow were monitored during 3 consecutive years in five *Acacia raddiana* trees and five *Acacia tortilis* trees in the Arid Arava Valley, southern Israel (annual precipitation 20–70 mm, restricted to October–May). We hypothesized that stem growth and other tree activities are synchronized with, and limited to single rainfall or flashflood events. Unexpectedly, cambial growth of both *Acacia* species was arrested during the wet season, and occurred during most of the dry season, coinciding with maximum daily temperatures as high as 45 °C and vapor pressure deficit of up to 9 kPa. Summer growth was correlated with peak sap flow in June, with almost year-round activity and foliage cover. To the best of our knowledge, these are the harshest drought conditions ever documented permitting cambial growth. These findings point to the possibility that summer cambial growth in *Acacia* under hyper-arid conditions relies on concurrent leaf gas exchange, which is in turn permitted by access to deep soil water. Soil water can support low-density tree populations despite heat and drought, as long as recharge is kept above a minimum threshold.

Keywords Acacia raddiana · Acacia tortilis · Leaf phenology · Sap flow · Desert · Global warming · Tree drought resistance · Arava

Communicated by Louis Stephen Santiago.

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s00442-018-4250-z) contains supplementary material, which is available to authorized users.

Tamir Klein tamir.klein@weizmann.ac.il

- ¹ The Dead Sea-Arava Science Center, Tamar Regional Council, 86910 Neve Zohar, Israel
- ² Department of Biological Sciences, Jaramogi Oginga Odinga University of Science and Technology, Bondo, Kenya
- ³ Institute of Soil, Water, and Environmental Sciences, Agricultural Research Organization, Rishon LeZion, Israel
- ⁴ Ecological Modelling, BayCEER, University of Bayreuth, Bayreuth, Germany
- ⁵ Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot, Israel

Introduction

Drought-induced tree mortality has been increasing in recent decades (Klein 2015; Anderegg et al. 2016), and is expected to increase further due to global warming and desertification, especially in already dry ecosystems (Steinkamp and Hickler 2015). Conversely, tree species which already survive under arid conditions could provide vital information on successful drought resistance strategies (Schwinning and Ehleringer 2001). Indeed, there is growing interest to elucidate some of these strategies, as demonstrated by recent studies on Tamarix and Populus in the Taklamakan desert (Gries et al. 2003; Lang et al. 2016) and Prosopis in the Atacama desert (Garrido et al. 2016). Water scarcity and high temperatures often interact to exacerbate drought stress on tree physiology (Ruehr et al. 2016). Hence, resolving tree drought resistance strategies, and specifically in arid environments, is a major interest of scientists and stakeholders alike (Hartmann et al. 2015; Cobb et al. 2018). In deserts, trees such as Acacia,

Tamarix, Populus, and *Prosopis* are often the only woody species, and as such, they are considered keystone species (Ward and Rohner 1997). Successful desert trees are essential locally for their associated ecosystems and important globally for their unique eco-physiology.

Among living tree species, Acacia raddiana and Acacia *tortilis* (the genus name changed to *Vachellia*¹) inhabit some of the hottest and driest places on Earth. These Acacia trees are major components of savannas and open woodlands in many arid regions of Africa and the Middle East (Maslin et al. 2003). Within the arid Arava valley, along the Syrian-African transform (Great Rift valley) in southern Israel and Jordan, A. raddiana and A. tortilis are the two most abundant and, in many places, the only tree species present (Danin 1983). In these arid habitats, Acacia trees are found mostly growing in the channels of ephemeral river beds ("wadis", a term from Arabic; Ward et al. 1993; Munzbergova and Ward 2002). Here, both A. raddiana and A. tortilis are considered keystone species that support the majority of the biodiversity surrounding them and locally improve soil conditions for other plant species (Milton and Dean 1995; Ward and Rohner 1997; Munzbergova and Ward 2002).

Although Acacia species dominate many of Earth's hottest and driest deserts, little is known about their growth dynamics in situ. Some knowledge of related eco-physiological parameters does exist but not for Acacia trees growing in situ and in arid environments. Water relations were studied extensively for A. tortilis, but not in hyper-arid conditions; for example, Do et al. (2008) studied water-use of A. tortilis in the northern Sahel, where annual rainfall ranged between 146 and 367 mm. Ludwig et al. (2003) studied hydraulic lift in A. tortilis trees on an East African savanna where the climate is tropical seasonal $(650 \pm 272 \text{ mm year}^{-1})$. Studies by Otieno et al. (2003, 2005a) were based on seeds that were collected in Kenya but brought back to Germany where they were germinated and grown under controlled conditions. Both Otieno et al. (2005b) and Do et al. (2008) showed the high drought resistance of this species, including physiological and morphological adaptations, in semi-arid environments, which do not represent the real extreme environments that some Acacias can tolerate. In the more arid Eastern Desert of Egypt (North Africa), wood anatomy of A. raddiana was studied using ¹⁴C, due to the lack of annual tree-rings (Andersen and Krzywinski 2007). In the Arava valley of Southern Israel, A. raddiana cambial activity was studied in seedlings and in shoots of mature trees (Fahn et al. 1968; Arzee et al. 1970). Several other studies have tried to address the challenging task of identifying water source(s) sustaining Acacia populations in the Arava. This topic has drawn particular attention, especially following mortality events due to consecutive drought years in the 1980s (Peled 1988; Shrestha et al. 2003). Based on the dynamics of A. raddiana shoot water potential across sites and seasons, it was suggested that surface floods are the major water source for populations in the Arava (Shrestha et al. 2003). Such a strategy is typical of pulse-driven arid ecosystems (Schwinning and Ehleringer 2001). Studying the water source for Acacia trees in situ has been done by exposing their root systems (Peled 1988), comparing the ${}^{18}O/{}^{16}O$ isotopic ratios in water samples extracted from Acacia twigs and from nearby water sources (Sher et al. 2010), and using Electrical Resistivity Tomography (ERT; Winters et al. 2015). Yet, none of these studies monitored in situ Acacia growth and water-use in stems and leaves to observe their temporal dynamics and relationships to local meteorological conditions.

Here, we aimed to decipher the eco-physiological dynamics of both *A. raddiana* and *A. tortilis*, the dominant tree species in the hyper-arid regions of North Africa and the Middle East, as a case of tree growth at the hot and dry extreme. We hypothesized that stem growth and other tree activities are synchronized with rainfall or flashflood events. We took advantage of state-of-the-art monitoring technologies, and used a high-resolution, long-term, and continuous observation approach. Electronic dendrometers, sap flow sensors, and phenology cameras were connected to individual trees, to produce an accurate and detailed representation of *Acacia* survival and growth in one of Earth's most hostile environments.

Materials and methods

Site and climate

Our study was conducted in Wadi Sheizaf, a dry sandy streambed at the northern edge of the Arid Arava Valley, southern Israel (Fig S1; 30°44'N, 35°14'E; elevation 137 below sea level). Meteorological data were obtained from the Israeli Meteorological Service for station 340528 in Hatzeva, located 7 km north of Wadi Sheizaf. The climate here is hot and dry: 30-year average minimum, mean, and maximum temperature of the hottest month were 26.2 °C, 33.2 °C, and 40.2 °C, respectively; average minimum, mean, and maximum temperature of the coolest month was recorded

¹ The genus (*Acacia*) was recently split (Kyalangalilwa et al. 2013) into two different genera, *Vachellia* and *Acacia*. While the original name (*Acacia* = "thorn" in latin) was reserved for the Australian (thornless) *Acacia* trees, the new name, *Vachellia*, was reserved for the *Acacias* from the rest of the world, with thorns. Searching the web science for scientifc papers published from 2015 to June 2016 showed that only 2.6% from the 346 Acacia papers published during this periord chose to use the new name. A short survey among Israel's botanists also demsonstrates that it is preferable to continue to use the genera name *Acacia* (see for example Winters et al. 2015; Nothers et al. 2017; Rodger et al. 2018). For these reasons, we chose to use the genera *Acacia*.

Fig. 1 Meteorological conditions at Hatzeva, 7 km north of Wadi Sheizaf (Arava valley, Israel; Fig. S1) in 2014–2016. During the 3 observation years, there were 77 days when temperature was > 40 °C, 26 days when vapor pressure deficit (VPD) was > 7 kPa, and only 4 days when rainfall was > 10 mm



as 9.1 °C, 14.4 °C, and 19.6 °C, respectively, and annual precipitation of only 20–70 mm is restricted to the period between October and May (Fig. 1). Multiple individual trees of *A. raddiana* and *A. tortilis* are scattered throughout the wadi, but never forming a continuous canopy (Fig. S1).

Stem growth

Stem circumference was measured continuously with automatic high-resolution radial dendrometers (DRL26, EMS, Brno, Czech Republic; Urban et al. 2013). These dendrometers measured circumference variations of less than 1 µm and logged the data into a built-in memory once per hour. In October-November 2013, dendrometers were installed on four A. raddiana trees, followed by two A. tortilis trees in April-June 2014. Later, in 2014, this setup was increased to five A. raddiana trees and six A. tortilis trees. Basal stem circumferences of the five A. raddiana trees were 111 cm, 129 cm, 47 cm, 104 cm, and 165 cm, and those of the six A. tortilis trees were 38 cm, 52 cm, 38 cm, 34 cm, 44 cm, and 51 cm. To ensure that circumference variations were not affected by the frequent temperature and radiation gradients, a 12th dendrometer was installed on a dead A. raddiana tree in December 2015. This dendrometer showed minimal changes, confirming the robustness of the dendrometer sensor. In the data analysis stage, all dendrometer values were set to zero on January 1st of each year. Tree-specific dendrometer data were used in four types of analyses: (1) diurnal stem growth (mm) during a given year, produced by zeroing readings on January 1st of each year; (2) relative stem growth (between 0 and 1), produced by dividing the values in (1) by the tree-specific total annual growth; (3) stem growth rate (mm day⁻¹), produced by subtracting the values in (1) from the values of the following day; (4) diurnal stem circumference variations (mm), produced by plotting the readings at hourly temporal resolution, and zeroing at midnight of each day.

Sap flow

Sap flow has been monitored continuously in six of the study trees (two A. raddiana and four A. tortilis trees) since November 2015. Thermal dissipation sensors (n=6) prepared in workshops at the University of Bayreuth (Germany) and at ARO Volcani center (Israel) were installed in the lower 30 cm of the stem, below the lowest branches, in the outer 2 cm of xylem, and connected to data loggers (CR1000, Campbell Scientific, UT, USA; and DL2e, Delta-T Devices, Cambridge, UK) and constant current power supplies. Probes were insulated by filling aluminum foil wraps tied around the stems and probes with polyurethane foam. Power was from solar charged 12 V car batteries. Sap flow measurement and analysis followed Granier and Loustau (1994). Stem circumference, which averaged 44 cm, was used to calculate sapwood area for each tree, assuming a linear decrease in flux density with depth, as described in Paudel et al. (2013) and Kanety (2014). We estimated zero flux to be 5 cm into the xylem, as found for many broadleaf, diffuse-porous tree species (Cohen et al. 2008). Sap flow $(L h^{-1})$ was calculated assuming zero flux at times of maximum temperature differentials measured during the 24 h period, which was typically found predawn. No further calibration factor was applied.

Foliage dynamics

To follow changes in tree phenology, all study trees (5 *A*. *raddiana* and 6 *A*. *tortilis*) were photographed monthly in



Fig. 2 Acacia stem growth in Wadi Sheizaf (Arava valley, Israel) in 2014–2016, **a–c**, respectively. Missing or truncated curves are due to measurement failures. Inset figure in **b** shows a zoom in on *A. tortilis* 102 during 15 July–15 August 2014 (see rectangle on the seasonal

curve of summer 2014). Most of the cambial growth occurred in May–June and August–November, coinciding with some of the hot-test and driest days

Wadi Sheizaf under natural irradiances. Photographs were taken between January 2016 and December 2016 from a marked spot and at consistent height and angle (from south to north, using the digital compass within the camera) and using a single camera (Nikon AW110, Japan). Photographs included a Kodak grey scale in the image, which was used for testing the effect of different time of day/season on color intensity.

Photographs were analyzed in R (R core Team 2013) using the recently described Phenopix (Filippa et al. 2016), an R package that allows for the extraction of phenological information from time-lapse digital photography of vegetation cover. A region of interest (ROI) was drawn onto an imported image and used to extract Red (R), Green (G), and Blue (B) (RGB) digital numbers (DN-range between 0 and 254), which allowed us to depict the greenness index (intensity of the Green channel/Red+Green+Blue) represented by the Green Chromatic Coordinate (GCC-range between 0 and 1). The ROI in Phenopix (Filippa et al. 2016) represented vector locations allowing for images that were taken from the same exact location/distance and angle to be automatically assessed for the same ROI, allowing us to process hundreds of images within minutes. To ensure that the dynamics of GCC of the tress showed real changes, we calculated the GCC of the grey patch [grey 182 (R = G = B = 182)] of the Kodak grey scale (Fig. S2). As the grey scale was placed at different locations of the images, the analysis was performed manually in ImageJ. To analyze the seasonal dynamics of the GCC, we fitted a generalized additive model (GAM) (Hastie and Tibshirani 1990). A GAM is a non-linear additive model that describes the relationship between the predictor and the response variable as a smooth function. We fitted different smoothing functions per species and allowed for random differences between trees.

Results

Acacia stem growth dynamics

Annual increments in Acacia stem circumference were between 0 and 40 mm, with large variations between individual trees and different years (Fig. 2). Differences between individual trees were not related to stem diameter (correlation $r^2 < 0.01$), i.e., trees with large annual increments included both larger and smaller individuals. There were no clear differences in stem growth between A. raddiana and A. tortilis trees. Stem growth in 2015 was smaller than in 2014 and 2016 (Fig. 2). In 2014 specifically, four of eight monitored trees grew > 30 mm, an annual increment that did not recur in the consecutive years. Stem growth was highly seasonal, with fast increments during May-June and again in August-November, and close to zero growth in December-April and in July. Interestingly, growth was decoupled from rainfall or flashflood events (Fig. 1), which are also restricted to November–April. Instead, trees were actively growing mostly when daily maximum temperature was > 27 °C, and up to 45 °C (June 2016). Except for the growth arrest in July, most of the annual stem increment was completed during such hot and dry days, with vapor pressure deficit (VPD) of up to 9 kPa, a combination of the very high air temperature and a very low relative humidity value of 6%. Except for a few days per year, stem circumference changes reflected growth per se and not elastic water-induced changes, typically identified by stem swelling and shrinking. For example, A. tortilis 102 showed a 0.2 mm decrease during late July 2014 (Fig. 2, inset), which was in turn dwarfed by a continuous increase of 22 mm during the following months.



Fig. 3 Cumulative relative rainfall amount and *Acacia* stem growth in Wadi Sheizaf (Arava valley, Israel) in 2014–2016, **a**–**c**, respectively. Missing or truncated curves are due to measurement failures. Curves

represent the fraction of the annual total stem increment per tree or rainfall amount

To test the sensitivity of stem growth to rainfall, relative growth curves and the cumulative rainfall amount (fractions of the annual totals) were superimposed (Fig. 3). In 2014 and 2015, stem growth took place mostly in spring and autumn, months after the majority of the rainfall amount had already been precipitated. However, in 2016, stem growth onset in many trees seemed to synchronize with the rain events (Fig. 3c). It is not clear whether the apparent cooccurrence in 2016 was the result of the delayed rains, or the earlier growth, compared to the two preceding years. In most trees and across the years, stem growth onset was usually not before 40% of the annual rain water input. However, overall, precipitation patterns alone could not sufficiently explain stem growth trends, such as the growth enhancement in August 2014 in most trees and the lack of growth before autumn 2015 despite the winter rains. Interestingly, the studied trees showed almost no stem shrinkage, which is often observed in dendrometry during periods of sub-optimal conditions (Chan et al. 2016; Zweifel et al. 2016). On an hourly scale, the diurnal stem circumference variations showed the expected daytime shrinking and nighttime swelling (Fig. S3), reflecting the changes in plant tissue water potential during the day (Steppe et al. 2015). The diurnal amplitude was ~ 0.3 mm during the growing season and < 0.1 mm outside the season (Fig. S3). A. raddiana had slightly larger amplitudes than A. tortilis.

The diurnal stem circumference increment was typically between 0 and 0.4 mm (Fig. 4). Short-term spikes of up to 1 mm day⁻¹ were mostly related to tissue swelling during rain events, and, hence, do not represent growth per se. In each of the 3 study years, there were roughly two growth periods, separated by two no-growth periods, in peak winter and peak summer, respectively. 2014 had the highest growth rates, but 2016 had the longest growth season, stopping only for periods of 5–20 days in February, April, August, and December (Fig. 4c). Summer growth was recorded in all 3 years in June, in July 2016, and in August 2014 and 2016. Among species, growth rates were mostly higher in *A. tortilis* in 2015, and in *A. raddiana* in 2016.

Plotting stem growth rate against meteorological conditions yielded no clear relationships, and hence, correlations were not significant (Fig. 5). Yet, it was clear that: (1) growth was not higher during rainy days; (2) most *A. raddiana* growth peaks followed rainy days, but were not related to rainfall amount; (3) growth of *A. raddiana* and *A. tortilis* was generally higher around 33 °C and 35 °C and around VPD of 3.8 kPa and 4.0 kPa, respectively, with local peaks in *A. raddiana* at lower temperature and VPD values.

Acacia sap flow dynamics

There was relatively little seasonal variability in the diurnal sap flow amounts (Fig. 6), and sap flow rates were $2-3 \text{ L h}^{-1}$ during mid-day hours on most days. The diurnal sap flux was around 12 L day⁻¹ year-round, increasing to over 13 L day⁻¹ in May–June and November, and decreasing to 7.4 L day⁻¹ at the end August (Fig. 6, inset). A morning spike reflecting high sap flow rates following sunrise was evident in most days and was up to 10 L h⁻¹ on cooler days, i.e., during November–February. The highest diurnal water-use was recorded in October–November (Fig. 6), and the second highest in May–June. Evidently, the highest sap flow activity coincided with the hottest and driest times, whereas lower activity was recorded on cooler, milder days, such as in March.



Date (Month-Year)

Fig.4 Acacia stem growth rate and cumulative relative rainfall amount in Wadi Sheizaf (Arava valley, Israel) in 2014–2016, **a–c**, respectively. Missing or truncated curves are due to measurement failures. The dashed horizontal lines at 0.05 and 0.5 mm day⁻¹ were

used to filter out periods of no-growth and rain-driven tissue swelling, respectively, in estimating growth periods (black bars in the bottom). A growth period was defined when at least two individual trees were growing at a rate > 0.05 mm day^{-1} (but < 0.5 mm day^{-1})



Fig. 5 Relationships between Acacia stem growth rate and meteorological conditions in Wadi Sheizaf (Arava valley, Israel) in 2014–2016. Each data point is a mean of 5 (6) trees of A. raddiana (A. tortilis)

Acacia leaf phenology

Crown foliage was maintained almost all year-round in both *A. raddiana* and *A. tortilis* (Fig. 7; Supplementary

Information Video S1). In the two species, new leaves emerged two times a year, in early March, and again in late October. The leaf shedding period was relatively short and timed to July (*A. raddiana*) or August (*A. tortilis*). In



Fig. 6 Diurnal courses of *A. tortilis* sap flow, air temperature (°C) and vapor pressure deficit (VPD in kPa) in Wadi Sheizaf (Arava valley, Israel) during 11 days in February–December 2016 (n=6). An annual curve with 10-day average±SE is in the inset. Sap flow morning

spikes occurred on cooler days, but the activity was roughly similar year-round, with the largest water-use at the hottest and driest days, in June



Fig.7 Leaf phenology of *A. raddiana* (top) and *A. tortilis* (bottom) in Wadi Sheizaf (Arava valley, Israel). Monthly photographs (shown here are examples of for *A. raddiana* 281 and *A. tortilis* 24) from six trees of each species were analyzed for the calculation of the greenness index represented by Green Chromatic Coordinate (GCC; dia-

grams on right-hand side). Both species had close to evergreen leaf habit, with two annual flushing events (early March and late October) and a short shedding period in July (*A. raddiana*; **a**) or August (*A. tortilis*; **b**) (color figure online)

general, seasonal changes in the foliage, and the resulting dynamics in greenness [Green Chromatic Coordinate (GCC)] were more pronounced in *A. tortilis* than *A. raddiana* (Fig. 7). The GCC changed between 0.36 (spring and autumn maxima points) and 0.33 (summer minima) in *A. raddiana*, and between 0.37 and 0.32 in *A. tortilis*. Leaf flushing was unrelated to water inputs, since the early March flushing preceded the two major rain events in late March and early April 2016, and the late October flushing coincided with only a 3 mm rain event (Fig. 1). The GCC of the grey scale (grey 182) hardly showed any seasonal effect (Fig. S2). This is in accordance with other studies that reported that GCC was relatively robust to variations in illumination (e.g., Toomey et al. 2015).

Discussion

Acacia year-round activity and summer growth

This study explored tree eco-physiology at the hot and dry edge of tree existence. The Arava valley in Israel offers some of the hottest and driest conditions on the planet, while still permitting Acacia tree year-round growth. The local climate is also characterized by large seasonal differences, with daily maximum temperatures spanning over an amplitude of 30 °C (Fig. 1) and large year-to-year variations in the amount and the timing of local rain events. In spite of these large variations, there was year-round growth activity in these trees. The diurnal sap flow curves showed robust water-use by the trees throughout the year (Fig. 6), and the crown foliage cover was close to evergreen, especially in A. raddiana (Fig. 7). On top of this remarkable persistence in a hostile environment, stem growth was unexpectedly timed to the hotter and drier months of the year (Figs. 2, 4). Therefore, our hypothesis of synchronization with single rainfall or flashflood events, also relying on the previous observations at the site (Shrestha et al. 2003), was rejected.

The early studies on wood anatomy of *A. raddiana* have already shown that cambial activity in twigs peaked in May and persisted until September, with very little growth in the cooler months (Fahn et al. 1968; Arzee et al. 1970). Using anatomical methods combined with ¹⁴C labelling, it was shown that May and August were the only months when both the cambium and phellogen were active (Fahn et al. 1968; Arzee et al. 1970). Our ultra-sensitive dendrometers confirm these previous observations in twigs, albeit in mature trees in the field, with continuous dynamics over 3 consecutive years.

The *Acacia* species studied here are of tropical origin and, hence, display thermophilic behavior (Otieno et al. 2005a, b). Indeed, cambial growth in our study was restricted to days with daily maximum temperature > 27 °C. Among the two species, the more drought-resistant A. tortilis (Otieno et al. 2003) grew faster than A. raddiana during the harsher 2015 (and slower in the following, better, year). A. tortilis growth rate peaked around 35 °C and was unrelated to rainfall (Fig. 5a, b). Why then was growth temporarily arrested in most trees during July or August? This was not related to harsher conditions in either of the months than in June, i.e., there was no temperature or VPD threshold limiting growth during these months. One possible explanation can be found in leaf phenology: leaf shedding and minimum greenness were observed in late July in A. raddiana and in early August in A. tortilis (Fig. 7; Video S1). The seasonal minimum in greenness corresponded with that of sap flow (Fig. 6, inset). If cambial growth relies on concurrent photosynthesis (see below), then leaf phenology has a direct limitation on growth. Defoliation decreases growth as well as tissue carbon reserves (Landhäusser and Lieffers 2012). Leaf shedding in high summer might be related to cumulative photo-damage, as shown in the pine Pinus halepensis growing on the edge of the Arava, 75 km north of Wadi Sheizaf (Yatir forest, 650 m above sea level; Maseyk et al. 2008). The global solar radiation in our site reaches over 1000 W m⁻² at noon in summer, and irreversible damage to the fully exposed photosynthetic tissue is probably unavoidable (Wittenberg et al. 2014). Yet, the controls of stem growth phenology seem to be more complex. For example, some A. tortilis trees were growing during September, in spite of the low foliage cover.

Inter-annual growth variations

Stem growth in 2015 was lower than in 2016, which was in turn lower than in 2014 (Fig. 2). Our analysis yielded limited sensitivity of stem growth to rain events (Fig. 3), with other factors such as temperature being involved. Yet, sensitivity to climate might act at longer time-scales. Summing the winter rainfall amounts preceding these summer seasons shows a rather similar inter-annual water input, with 69 mm, 66 mm, and 58 mm in 2014, 2015, and 2016, respectively. In general, inter-annual variation in precipitation in the Arava can be much larger (Ginat et al. 2011). Why then was growth so different between the years? In arid environments, the total rainfall amount might be less important than the temporal rainfall distribution (Raz-Yaseef et al. 2010a, b; Klein et al. 2014) or the spatial distribution (Isaacson et al. 2017). Our measurements are in agreement with the previous observations at the site, showing higher tree water stress following a dry winter than during the dry, hot summer (Shrestha et al. 2003). The ranking of growth by years 2014 > 2016 > 2015 correlates with two such observations: (1) the last substantial rain event ("spring rain") was 7th May 2014 > 12th April 2016 > 28th March 2015. This entails over a month of inter-annual differences in soil water conditions;

(2) the number of days with rainfall > 10 mm was 3 > 1 > 0 in 2014, 2016, and 2015, respectively. This means higher percolation and lower losses to surface evaporation in 2014. The role of indirect water inputs must be considered too. Belowground flows can feed an aquifer which the *Acacia* roots may have access to (Winters et al. 2015), but this is yet to be studied.

Carbon source for growth

Can concurrent photosynthesis explain the summer cambial growth? Active carbon uptake at temperature > 40 °C and VPD>8 kPa is hard to imagine. Trees typically cease growth at temperature >40 °C and VPD >6 kPa (Ruehr et al. 2016). However, in our trees, leaves were there almost all year-round and the trees maintained relatively moderate and stable sap flow rates during the entire experimental period. The desert environment causes large seasonal amplitudes (Fig. 1), as well as large diurnal amplitudes. For example, VPD < 2 kPa was recorded almost every night throughout the year, including summer (Fig. 6). In a typical day in May 2016, VPD was 1.74 kPa and 2.61 kPa at 8:00 and 11:00, respectively. In addition, in a typical day in August 2016, VPD was 1.93 kPa and 2.90 kPa in 8:00 and 11:00, respectively. These VPD levels usually permit some stomatal conductance, and hence, photosynthetic activity could take place for 4 h or more every morning (Klein et al. 2016). It is possible that during a summer day stomata shut down in late morning, while sap flow continues until sunset (Fig. 6), to refill water storage in woody or leaf tissues, or even embolisms in the xylem (Klein et al. 2016). Such physiological regulation can be regarded as drought avoidance at a fine temporal scale.

Water source for growth

Our sap flow measurements suggest a rich and constant year-round water source for Acacia growing in the desert. Studying the roles and interactions of soil water, ground water, and Acacia root accessibility was beyond the scope of this current research. Nevertheless, we have indications of significant water amounts at 7-10 m depth in a nearby site (7 km to the north; Winters et al. 2015), and their use by A. raddiana (Sher et al. 2010). Accessible ground water support Prosopis trees even in the Atacama, the driest desert on Earth, as well as Tamarix and Populus in the Taklamakan, with water table depth playing a major role in tree function (Garrido et al. 2016; Gries et al. 2003, respectively). Desert trees have the deepest roots in the plant kingdom (Canadell et al. 1996), and Acacia species rank high among them, e.g., records of A. raddiana roots at 35 m below surface (Stone and Kalisz 1991). In the Mojave Desert, deep-rooted plants were less vulnerable to drought than others on soils that allowed for deep-water percolation (Munson et al. 2015). *Acacia* roots go deep, as well as far from the stem, to collect water from a large area as possible. In a road construction site in the Arava valley, 120 km south of Wadi Sheizaf, there was an attempt to conserve *A. tortilis* trees by digging around trees, leaving them on 2–3 m wide, 7–8 m tall poles (Fig. S4). Failure of this attempt and the resulting *Acacia* tree die-off indicates the significant role of lateral roots for their survival.

The observed inter-annual variation in growth and its relation to rainfall distribution (Fig. 2) indicates that the water source for these trees, as deep as it may be, is also sensitive to climate. Soil water can support low-density tree populations despite heat and drought, as long as recharge is kept above a minimum, yet unknown, threshold. Furthermore, the variation among individual trees in 2014 suggests different access to water, either due to source patchiness (water pockets), or root system architecture. Hydrological niche segregation is a widespread phenomenon across terrestrial biomes, but, so far, only arid plant communities have demonstrated temporal partitioning that promotes coexistence (Silvertown et al. 2015). Acacia trees, typically growing in savanna ecosystems, are used to competition with grass species over water resources (Cramer et al. 2010). In that sense, desert ecosystems, with only limited grass growth, offer an advantage for Acacia growth.

In pulse-driven arid ecosystems, four major plant forms have evolved: winter annuals, drought-deciduous perennials, evergreen perennials, and stem succulents (Schwinning and Ehleringer 2001). The third category fits best the *Acacia* studied here, and predicts large root/shoot ratio, deep root system, and lower leaf conductance with low stomatal sensitivity, as adaptations that maximize deeper soil water-use (Schwinning and Ehleringer 2001). In the case of *Acacia* in the hyper-arid Arava, while some indications exist for deep soil water-use (Sher et al. 2010; Winters et al. 2015), predictions for the other traits should be used as major hypotheses for future research at our site.

Implications for arid ecosystems under climate change

In face of climate change and its threats to forests worldwide, our study points to the potential advantage of lowdensity woodland over a forest, allowing each individual tree to exploit a large area by growing proliferate root systems. Under future drying and warming scenarios, such sparse woodlands might survive climate change better than dense forests. On an ecological perspective, we demonstrate that in spite of flashflood and rainfall pulses, deep soil water in our hyper-arid ecosystem had a larger effect on *Acacia* growth. This also indicates that terrestrial vegetation models must account for belowground conditions to capture ecosystem dynamics. In addition, future research should quantify the hydrological threshold for *Acacia* growth and survival, to facilitate better predictions under climate change.

Acknowledgements We thank Victor Lukyanov (ARO Volcani Center, Israel) for technical assistance in the field. TK wishes to thank the Benoziyo Fund for the Advancement of Science; Mr. and Mrs. Norman Reiser, together with the Weizmann Center for New Scientists; the Edith and Nathan Goldberg Career Development Chair. CB and DO thank the Bayrische Forschungs (BayFor) Alianz, Germany, for supporting the visits to the Arava. GW thanks the Arava Drainage Authority and the Israeli Ministry of Science and Technology (MOST) for their continued support. Thanks to the anonymous reviewers and to the editor, Russel K. Monson, for their helpful comments and suggestions which significantly improved the MS.

Author contribution statement GW initiated the study in 2013, with monitoring operated together with GR. DO, SC, and TK joined the study in 2015, with YW, IR, IP, and CB joining in 2016. TK started the analysis and manuscript drafting, with contributions from all authors.

Data accessibility Data used in this paper will be archived in Figshare.

Compliance with ethical standards

Conflict of interest The authors declare no conflict of interest.

References

- Anderegg WR, Klein T, Bartlett M, Sack L, Pellegrini AF, Choat B, Jansen S (2016) Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. Proc Natl Acad Sci 113:5024–5029
- Andersen GL, Krzywinski K (2007) Longevity and growth of Acacia tortilis; insights from ¹⁴C content and anatomy of wood. BMC Ecol 7:4
- Arzee T, Waisel Y, Liphschitz N (1970) Periderm development and phellogen activity in the shoots of *Acacia raddiana* Savi. New Phytol 69(2):395–398
- Canadell J, Jackson RB, Ehleringer JB, Mooney HA, Sala OE, Schulze ED (1996) Maximum rooting depth of vegetation types at the global scale. Oecologia 108(4):583–595
- Chan T, Hölttä T, Berninger F, Mäkinen H, Nöjd P, Mencuccini M, Nikinmaa E (2016) Separating water-potential induced swelling and shrinking from measured radial stem variations reveals a cambial growth and osmotic concentration signal. Plant Cell Environ 39(2):233–244
- Cobb R et al (2018) Ecosystem dynamics and management after forest die-off: a global synthesis with conceptual state-and-transition models. Ecosphere 8(12):e02034
- Cohen Y, Cohen S, Cantuarias-Aviles T, Schiller G (2008) Variations in the radial gradient of sap velocity in trunks of forest and fruit trees. Plant Soil 305(1):49–59
- Cramer MD, Van Cauter A, Bond WJ (2010) Growth of N2-fixing African savanna *Acacia* species is constrained by below-ground competition with grass. J Ecol 98:156–167
- Danin A (1983) Desert vegetation of Israel and Sinai. Cana Publishing House, Jerusalem
- Do FC, Rocheteau A, Diagne AL, Goudiaby V, Granier A, Lhomme JP (2008) Stable annual pattern of water use by *Acacia tortilis* in Sahelian Africa. Tree Physiol 28(1):95–104

- Fahn A, Waisel Y, Benjamini L (1968) Cambial activity in Acacia raddiana Savi. Ann Bot 32(3):677–686
- Filippa G et al (2016) Phenopix: a R package for image-based vegetation phenology. Agric For Meteorol 220:141–150
- Garrido M, Silva P, Acevedo E (2016) Water relations and foliar isotopic composition of *Prosopis tamarugo* Phil., an endemic tree of the Atacama desert growing at three levels of water table depth. Plant Sci 7:375
- Ginat H, Shlomi Y, Batarseh S, Vogel J (2011) Reduction in precipitation levels in the Arava Valley (southern Israel and Jordan), 1949–2009. J Dead Sea Arava Res 1:1–7
- Granier A, Loustau D (1994) Measuring and modelling the transpiration of a maritime pine canopy from sap-flow data. Agric For Meteorol 71(1):61–81
- Gries D, Zeng F, Foetzki A, Arndt SK, Bruelheide H, Thomas FM, Zhang X, Runge M (2003) Growth and water relations of *Tamarix ramosissima* and *Populus euphratica* on Taklamakan desert dunes in relation to depth to a permanent water table. Plant Cell Environ 26:725–736
- Hartmann H, McDowell NG, Trumbore S (2015) Allocation to carbon storage pools in Norway spruce saplings under drought and low CO₂. Tree Physiol 35(3):243–252
- Hastie T, Tibshirani RJ (1990) Generalized additive models. Chapman and Hall/CRC, Boca Raton
- Isaacson S, Ephrath JE, Rachmilevitch S, Maman S, Ginat H, Blumberg DG (2017) Long and short term population dynamics of Acacia trees via remote sensing and spatial analysis: case study in the southern Negev Desert. Remote Sens Environ 198:95–104
- Kanety T, Naor A, Gips A, Dicken U, Lemcoff JH, Cohen S (2014) Irrigation influences on growth, yield, and water use of persimmon trees. Irrig sci 32(1):1–13
- Klein T (2015) Drought-induced tree mortality: from discrete observations to comprehensive research. Tree Physiol 35(3):225–228
- Klein T, Rotenberg E, Cohen-Hilaleh E, Raz-Yaseef N, Tatarinov F, Preisler Y, Ogée J, Cohen S, Yakir D (2014) Quantifying transpirable soil water and its relations to tree water use dynamics in a water-limited pine forest. Ecohydrology 7:409–419
- Klein T, Cohen S, Paudel I, Preisler Y, Rotenberg E, Yakir D (2016) Diurnal dynamics of water transport, storage and hydraulic conductivity in pine trees under seasonal drought. iForest 9:710–719
- Kyalangalilwa B, Boatwright JS, Daru BH, Maurin O, Der Van, Bank M (2013) Phylogenetic position and revised classification of *Acacia* s.l. (Fabaceae: Mimosoideae) in Africa, including new combinations in *Vachellia* and *Senegalia*. Bot J Linn Soc 172(4):500–523
- Landhäusser SM, Lieffers VJ (2012) Defoliation increases risk of carbon starvation in root systems of mature aspen. Trees 26(2):653–661
- Lang P, Ahlborn J, Schäfer P, Wommelsdorf T, Jeschke M, Zhang X, Thomas FM (2016) Growth and water use of *Populus euphratica* trees and stands with different water supply along the Tarim River, NW China. For Ecol Manag 380:139–148
- Ludwig F, Dawson TE, Kroon H, Berendse F, Prins HH (2003) Hydraulic lift in *Acacia tortilis* trees on an East African savanna. Oecologia 134(3):293–300
- Maseyk KS, Lin T, Rotenberg E, Grünzweig JM, Schwartz A, Yakir D (2008) Physiology–phenology interactions in a productive semiarid pine forest. New Phytol 178:603–616
- Maslin BR, Miller JT, Seigler DS (2003) Overview of the generic status of Acacia (Leguminosae: Mimosoideae). Aust Syst Bot 16(1):1–18
- Milton SJ, Dean WRJ (1995) How useful is the keystone species concept, and can it be applied to *Acacia erioloba* in the Kalahari Desert? J Nat Conserv 4(3):147–156
- Munson SM, Webb RH, Housman DC, Veblen KE, Nussear KE, Beever EA, Hartney KB, Miriti MN, Phillips SL, Fulton RE,

Tallent NG, Smith M (2015) Long-term plant responses to climate are moderated by biophysical attributes in a North American desert. J Ecol 103:657–668

- Munzbergova Z, Ward D (2002) Acacia trees as keystone species in Negev desert ecosystems. J Veg Sci 13(2):227–236
- Nothers M, Segev N, Kreyling J, Hjazin A, Groner E (2017) Desert vegetation forty years after an oil spill. J environ qual 46(3):568–575
- Otieno DO, Kinyamario JI, Omenda TO (2003) Growth features of *Acacia tortilis* and *Acacia xanthophloea* seedlings and their response to cyclic soil drought stress. Afr J Ecol 39(2):126–132
- Otieno DO, Schmidt MWT, Adiku S, Tenhunen J (2005a) Physiological and morphological responses to water stress in two *Acacia* species from contrasting habitats. Tree Physiol 25(3):361–371
- Otieno DO, Schmidt MWT, Kinyamario JI, Tenhunen J (2005b) Responses of *Acacia tortilis* and *Acacia xanthophloea* to seasonal changes in soil water availability in the savanna region of Kenya. J Arid Environ 62(3):377–400
- Paudel I, Kanety T, Cohen S (2013) Inactive xylem can explain differences in calibration factors for thermal dissipation probe sap flow measurements. Tree Physiol 33(9):986–1001
- Peled Y (1988) The mortality of *Acacia* trees in the southern Arava region. Thesis submitted to the Hebrew Univ., Jerusalem (Israel). Faculty of Agriculture. Dept. of Horticulture
- R Core Team (2013) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. http:// www.R-project.org/. Accessed 9 Apr 2018
- Raz-Yaseef N, Rotenberg E, Yakir D (2010a) Effects of spatial variations in soil evaporation caused by tree shading on water flux partitioning in a semi-arid pine forest. Agric For Meteorol 150(3):454–462
- Raz-Yaseef N, Yakir D, Rotenberg E, Schiller G, Cohen S (2010b) Ecohydrology of a semi-arid forest: partitioning among water balance components and its implications for predicted precipitation changes. Ecohydrology 3:143–154
- Rodger YS, Greenbaum G, Silver M, Bar-David S, Winters G (2018) Detecting hierarchical levels of connectivity in a population of *Acacia tortilis* at the northern edge of the species' global distribution: combining classical population genetics and network analyses. PLoS One 13:e0194901
- Ruehr NK, Gast A, Weber C, Daub B, Arneth A (2016) Water availability as dominant control of heat stress responses in two contrasting tree species. Tree Physiol 36(2):164–178

- Schwinning S, Ehleringer JR (2001) Water use trade-offs and optimal adaptations to pulse-driven arid ecosystems. J Ecol 89:464–480
- Sher AA, Wiegand K, Ward D (2010) Do Acacia and Tamarix trees compete for water in the Negev desert? J Arid Environ 74(3):338–343
- Shrestha MK, Stock WD, Ward D, Golan-Goldhirsh A (2003) Water status of isolated Negev desert populations of *Acacia raddiana* with different mortality levels. Plant Ecol 168(2):297–307
- Silvertown J, Araya Y, Gowing D (2015) Hydrological niches in terrestrial plant communities: a review. J Ecol 103(1):93–108
- Steinkamp J, Hickler T (2015) Is drought-induced forest dieback globally increasing? J Ecol 103(1):31–43
- Steppe K, Sterck F, Deslauriers A (2015) Diel growth dynamics in tree stems: linking anatomy and ecophysiology. Trends Plant Sci 20(6):335–343
- Stone EL, Kalisz PJ (1991) On the maximum extent of tree roots. For Ecol Manag 46(1):59–102
- Toomey M, Friedl MA, Frolking S et al (2015) Greenness indices from digital cameras predict the timing and seasonal dynamics of canopy-scale photosynthesis. Ecol Appl 25(1):99–115
- Urban J, Bednářová E, Plichta R, Kučera J (2013) Linking phenological data to ecophysiology of European Beech. Acta Hortic 991:293–299
- Ward D, Rohner C (1997) Anthropogenic causes of high mortality and low recruitment in three Acacia tree taxa in the Negev desert, Israel. Biodivers Conserv 6(6):877–893
- Ward D, Olsvig-Whittaker L, Lawes M (1993) Vegetation-environment relationships in a Negev Desert erosion cirque. J Veg Sci 4(1):83–94
- Winters G, Ryvkin I, Rudkov T, Moreno Z, Furman A (2015) Mapping underground layers in the super arid Gidron Wadi using electrical resistivity tomography (ERT). J Arid Environ 121:79–83
- Wittenberg G, Levitan A, Klein T, Dangoor I, Keren N, Danon A (2014) Knockdown of the Arabidopsis thaliana chloroplast protein disulfide isomerase 6 results in reduced levels of photoinhibition and increased D1 synthesis in high light. Plant J 78(6):1003–1013
- Zweifel R, Haeni M, Buchmann N, Eugster W (2016) Are trees able to grow in periods of stem shrinkage? New Phytol 211(3):839–849